

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/00, 15/09, 15/12</b>	<b>A1</b>	(11) International Publication Number: <b>WO 95/29233</b> (43) International Publication Date: 2 November 1995 (02.11.95)
<p>(21) International Application Number: PCT/US95/04860</p> <p>(22) International Filing Date: 19 April 1995 (19.04.95)</p> <p>(30) Priority Data: 08/230,491 20 April 1994 (20.04.94) US</p> <p>(71) Applicant: LUDWIG INSTITUTE FOR CANCER RE- SEARCH [US/US]; 1345 Avenue of the Americas, New York, NY 10105 (US).</p> <p>(72) Inventors: RETTIG, Wolfgang, J.; Amriswilstrasse 7, D- 88400 Biberach (DE). SCANLAN, Matthew, J.; 1275 York Avenue, New York, NY 10021 (US). GARIN-CHESA, Pilar; Amriswilstrasse 7, D-88400 Biberach (DE). OLD, Lloyd, J.; 1345 Avenue of the Americas, New York, NY 10105 (US).</p> <p>(74) Agent: HANSON, Norman, D.; Felfe &amp; Lynch, 805 Third Avenuc, New York, NY 10022 (US).</p>		<p>(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: ISOLATED NUCLEIC ACID MOLECULE CODING FOR FIBROBLAST ACTIVATION PROTEIN <math>\alpha</math> AND USES THEREOF</p> <p>(57) Abstract</p> <p>The invention describes the identification and isolation of nucleic acid molecules which code for fibroblast activation protein alpha, or "FAP<math>\alpha</math>". Various applications of the isolated molecules are also described.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

ISOLATED NUCLEIC ACID MOLECULE CODING FOR FIBROBLAST  
ACTIVATION PROTEIN  $\alpha$  AND USES THEREOF

FIELD OF THE INVENTION

This invention relates to certain molecules associated with cancer cells and reactive with tumor stromal cells. More particularly, it relates to fibroblast activation protein alpha ("FAP $\alpha$ " hereafter). The molecule has previously been identified immunologically, but nucleic acid molecules coding for it had not been isolated or cloned. This, inter alia, is the subject of the invention. The protein has a molecular weight of from about 88 to about 95 kilodaltons as determined by SDS-PAGE. This molecule is characterized by a number of features and properties which are shared by and characteristic of membrane bound enzymes, suggesting very strongly that it, too, is a membrane bound enzyme. The nucleic acid molecules, which are a key part of the invention, are useful both as probes for cell expressing FAP $\alpha$ , and as starting materials for recombinant production of the protein. The recombinant protein can then be used to produce monoclonal antibodies specific for the protein, and are thus useful diagnostic agents themselves.

BACKGROUND AND PRIOR ART

The invasive growth of epithelial cancers is associated with characteristic cellular and molecular changes in the supporting stroma. For example, epithelial cancers induce the formation of tumor blood vessels, the recruitment of reactive tumor stromal fibroblasts, lymphoid and phagocytic infiltrates, the release of peptide mediators and proteolytic enzymes, and the production of an altered extracellular matrix (ECM). See, e.g., Folkman, Adv. Cancer Res. 43: 175-203 (1985); Basset et al., Nature 348: 699-704 (1990); Denekamp et al., Cancer Metastasis Rev. 9: 267-282 (1990); Cullen et al., Cancer Res. 51: 4978-4985 (1991); Dvorak et al., Cancer Cells 3: 77-85 (1991); Liotta et al., Cancer Res. 51: 5054s-5059s (1991); Garin-Chesa et al., J. Histochem. Cytochem. 37: 1767-1776 (1989). A highly consistent molecular trait of the stroma in several common histologic types of epithelial

5 cancers is induction of the fibroblast activation protein (FAP $\alpha$ ), a cell surface glycoprotein with an observed M<sub>r</sub> of 95,000 originally discovered with a monoclonal antibody, mAb F19, raised against proliferating cultured fibroblasts. See Rettig et al., Cancer Res. 46: 6406-6412 (1986); Rettig et al., Proc. Natl. Acad. Sci. USA 85: 3110-3114 (1988); Garin-Chesa et al., Proc. Natl. Acad. USA 87: 7235-7239 (1990); 10 Rettig et al., Cancer Res. 53: 3327-3335 (1993). Each of these four papers is incorporated by reference in its entirety.

15 Immunohistochemical studies such as those cited supra have shown that FAP $\alpha$  is transiently expressed in certain normal fetal mesenchymal tissues but that normal adult tissues are generally FAP $\alpha$ <sup>-</sup>. Similarly, malignant epithelial, neural and hematopoietic cells are FAP $\alpha$ <sup>-</sup>. However, most of the 20 common types of epithelial cancers, including >90% of breast, lung, skin, pancreas, and colorectal carcinomas, contain abundant FAP $\alpha$ <sup>+</sup> reactive stromal fibroblasts. Garin-Chesa et al., Proc. Natl. Acad. Sci. USA 87: 7235-7239 (1990). The FAP $\alpha$ <sup>+</sup> tumor stromal fibroblasts almost invariably accompany 25 newly-formed tumor blood vessels, forming a distinct cellular compartment interposed between the tumor capillary endothelium and the basal aspect of malignant epithelial cell clusters. While FAP $\alpha$ <sup>+</sup> stromal fibroblasts are found in both primary and metastatic carcinomas, benign and premalignant epithelial 30 lesions, such as fibroadenomas of the breast and colorectal adenomas only rarely contain FAP $\alpha$ <sup>+</sup> stromal cells. In contrast to the stroma-specific localization of FAP $\alpha$  in epithelial neoplasms, FAP $\alpha$  is expressed in the malignant cells of a large proportion of bone and soft tissue sarcomas. Rettig et al., Proc. Natl. Acad. Sci. USA 85: 3110-3114 (1988). Finally, 35 FAP $\alpha$ <sup>+</sup> fibroblasts have been detected in the granulation tissue of healing wounds (Garin-Chesa et al., supra). Based on the restricted distribution pattern of FAP $\alpha$  in normal tissues and its uniform expression in the supporting stroma of many 40 epithelial cancers, clinical trials with <sup>125</sup>I-labeled mAb F19 have been initiated in patients with metastatic colon cancer

5 (Welt et al., Proc. Am. Assoc. Cancer Res. 33: 319 (1992)) to explore the concept of "tumor stromal targeting" for immunodetection and immunotherapy of epithelial cancers.

The induction of FAP $\alpha$ <sup>+</sup> fibroblasts at times and sites of tissue remodeling during fetal development, tissue repair, and carcinogenesis is consistent with a fundamental role for this molecule in normal fibroblast physiology. Thus, it is of interest and value to isolate and to clone nucleic acid molecules which code for this molecule. This is one aspect of the invention, which is described in detail together with other features of the invention, in the disclosure which follows.

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows results obtained from immunoprecipitation studies carried out on detergent extracts of Trans <sup>35</sup>S-labeled cells. The study was designed to immunoprecipitate FAP $\alpha$  and CD26. Cell types were SW872, which is a human sarcoma cell line, COS-FAP, which is a cell line transfected with a vector coding for FAP $\alpha$ , i.e., pFAP-38, described in the application, and COS-CD26, which is a COS cell line transfected with a CD26 coding plasmid. Extracts were precipitated with anti-FAP $\alpha$  monoclonal antibody F19, anti-CD26 mAb EF-1, or a negative control mouse Ig.

Figure 2A presents Northern blot analysis of FAP $\alpha$  expression in a cell line (ovarian cancer SK-OV6), which has FAP $\alpha$ <sup>+</sup>/CD26<sup>+</sup> phenotype), as well as two cell lines (fibroblasts WI-38 and GM 05389), which have FAP $\alpha$ <sup>+</sup>/CD26<sup>+</sup> phenotype.

Figure 2B shows  $\gamma$ -actin expression for the cell lines of figure 2A.

Figure 3 compares the deduced amino acid sequence for FAP $\alpha$ , and the known sequence of CD26. The alignment has been optimized.

Figure 4 depicts heterodimer formation between FAP $\alpha$  and CD26 in COS-1 transfectants.

Figures 5A-5H, inclusive, display immunohistochemical detection of FAP $\alpha$  and CD26 in various cancers. In figures 5A and 5B, breast cancer is studied, for FAP $\alpha$  (figure 5A), and

5 CD26 (figure 5B). In figures 5C and 5D, malignant fibrous histiocyoma are studied, for FAP $\alpha$  (figure 5C), and CD26 (figure 5D). Dermal scar tissue is examined in figures 5E (FAP $\alpha$ ), and 5F (CD26). Renal cell carcinoma is studied in figure 5G (FAP $\alpha$ ), and 5H (CD26).

10 DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Example 1

Fibroblast cell line WI-38 had been observed, previously, to react with mAb F19 (Rettig et al., Canc. Res. 46: 6406-6412 (1986); Rettig et al., Proc. Natl. Acad. USA 85: 3110-3114 (1988); Garin-Chesa et al., Proc. Natl. Acad. Sci. USA 87: 7235-7239 (1990); Rettig et al., Canc. Res. 53: 3327-3335 (1993)). It was used in the experiments which follow.

20 A cDNA library was prepared from WI-38, using well known techniques and commercially available materials. Specifically, the library was constructed in expression vector pCDNAI, using the Fast Track mRNA isolation kit, and Librarian cDNA phagemid system. Once the library was prepared, the vectors were electroporated into cell line *E. coli* MC 1061/P3. The pCDNAI expression vector contains an antibiotic resistance gene, so the *E. coli* were selected via antibiotic resistance. 25 The colonies which were resistant were then used in further experiments. The plasmid DNA from the colonies was obtained via alkaline lysis and purification on CsCl<sub>2</sub>, in accordance with Sambrook et al, Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Lab, Cold Spring Harbor, N.Y. 2d Ed. 30 1989). The technique is well known to the art, but is incorporated by reference herein.

Once the plasmid DNA was isolated, it was used to transfect COS-1 cells, which were then cultured for forty- 35 eight hours, after which these was tested with antibody coated dishes. The mAbs used included F19, as described by Rettig et al., (1986), supra, which is incorporated by reference in its entirety. As COS-1 cells are normally FAP $\alpha$ <sup>-</sup>, any positive results indicated the presence of the coding sequence. The immunoselection protocol was that of Aruffo et al., Proc. 40 Natl. Acad. Sci USA 84: 3365-3369 (1987), incorporated by

5 reference herein.

Plasmid DNA from positive clones was recovered, in accordance with Hirt, J. Mol. Biol. 26: 365-369 (1967), reintroduced into E. coli MC 1061/P3, and reselected into COS-1 cells.

10 The protocol presented herein was followed for four rounds. After this, the plasmid DNA of 50 isolated bacterial colonies was purified, using the Qiagen plasmid kit. Of the colonies, 27 clones were found to contain identical 2.8 kb inserts, as determined by EcoRI restriction enzyme mapping.  
15 Several of these were found to contain FAP $\alpha$ -specific cDNA, via transient expression in COS-1 cells and direct immunofluorescence staining. One of these clones, i.e., "pFAP.38" was selected for further study, as elaborated upon infra.

20 Example 2

Once pFAP.38 had been identified, it was tested together with a vector coding for known cell surface marker CD26 ("pCD26"), as well as with control vector pCDNA I.

25 In these experiments, COS-1 cells were transfected with one of pFAP.38, pCD26, or pCDNAI. After forty-eight hours, the transfectants were tested, using the well known MHA rosetting assay for cell surface antigen expression. In these experiments, mAb F19, which is FAP $\alpha$  specific, was used, together with mAb EF-1, which is CD26 specific. Also used  
30 were four other FAP $\alpha$  specific mAbs, i.e., FB23, FB52, FB58 and C48. Also tested were two cancer cell lines, which are known to react with mAb F19 (SW872 liposarcoma), or EF-1 (SK-OV6 ovarian cancer). The results are set forth in Table 1, which follows.



5 Table 1. Cell surface expression of multiple FAP $\alpha$  epitopes and CD26 in human cells and COS-1 cell transfectants.

10	Target Cell	F19	FB23	FB52	FB58	C48	EF-1
	<u>Human cells</u>						
15	SW872 liposarcoma	95%	>95%	>95%	>95%	>95%	-
	SW-OV6 ovarian cancer	-	-	-	-	-	>95%
	<u>COS-1 transfectants</u>						
20	COS-pCDNAI control	-	-	-	-	-	-
	COS-pFAP 38	40%	30%	40%	20%	20%	-
25	COS-pCD26	-	-	-	-	-	40%

### Example 3

Immunoprecipitation studies were then carried out to identify the antigen being targeted by the antibodies.

30 Cells were metabolically labelled with Trans <sup>35</sup>S-label, (ICN), extracted with lysis buffer (0.01 M Tris-HCl/0.15 M NaCl/0.01 M MgCl<sub>2</sub>/0.5% Nonidet P-40/aprotinin (20 ug/ml)/2 mM phenylmethyl- sulfonyl fluoride), and then immunoprecipitated. The protocols used are all well known, as will be seen by reference to Rettig et al., Canc. Res. 53: 3327-3335 (1993);  
 35 and Fellingner et al., Canc. Res. 51: 336-340 (1991), the disclosures of which are all incorporated by reference in their entirety. Precipitating mAbs were negative control mouse Ig, mAb F19, or EF-1. Control tests were carried out  
 40 with mock transfected COS-1 cells. Following immunoprecipitation, the immunoprecipitates were separated on NaDodSO<sub>4</sub>/PAGE, under reducing conditions. In some experiments, an additional test was carried out to determine whether or not the immunoprecipitated material was  
 45 glycosylated. In these experiments, cell extracts were fractionated with Con A-SEPHAROSE prior to immunoprecipitation. Following immunoprecipitation, but prior to fractionation on NaDodSO<sub>4</sub>/PAGE, these precipitates were

5 digested with N-Glycanase.

The results are shown in figure 1. In COS-1 cells, pFAP.38 directs expression of an 88 kd protein species (as determined via SDS-PAGE), which is slightly smaller than the 95 kd FAP $\alpha$  species produced by SW872, or cultured fibroblasts. Digestion with N-Glycanase produced peptides of comparable size (i.e., 74 kd versus 75 kd), showing that the glycosylation of the FAP $\alpha$  protein in COS-1 cells is different than in the human cell lines.

#### Example 4

15 Classic Northern blot analysis was then carried out, using the mRNA from FAP $\alpha$  fibroblast cell lines WI-38 and GM 05389, and FAP $\alpha$  ovarian cancer cell line SK-OV6. Using the procedures of Sambrook et al., supra, five micrograms of mRNA from each cell line were tested. The probes used were <sup>32</sup>P  
20 labelled, and were prepared from a 2.3 kb ECO I fragment of pFAP.38, a 2.4 kb Hind III fragment of CD26, and a 1.8 kb BamHI fragment of  $\gamma$ -actin cDNA. These fragments had been purified from 1% agarose gels.

Figure 2 presents these results. The extracts of FAP $\alpha$  fibroblast strains show a 2.8 kb FAP mRNA species, but  
25 extracts of SK-OV6 do not. A  $\gamma$ -actin mRNA species (1.8 kb), is seen in all species.

#### Example 5

The cDNA identified as coding for FAP $\alpha$  was subjected to  
30 more detailed analysis, starting with sequencing. The classic Sanger methodology, as set forth in Proc. Natl. Acad. Sci. USA 74: 5463-5467 (1977), was used to sequence both strands of the cDNA. Once this was secured, an amino acid sequence was deduced therefrom. This information is presented in SEQ ID  
35 NO: 1. The sequence was then compared to the known amino acid sequence of CD26 (Morimoto et al., J. Immunol. 143: 3430-3437 (1989)). Figure 3 presents the comparison, using optimized sequence alignment. Any gaps in the comparison are indicated by asterisks, while identical amino acids are shown by dashes  
40 in the CD26 sequence. A hydrophobic, putative transmembrane sequence is double underlined, while potential N-glycosylation

5 sites are single underlined.

The sequence analysis shows a 2812 base pair insert, wherein 2277 base pairs constitute the open reading frame. This ORF extends from start codon ATG at nucleotide 209, to stop codon TAA at 2486.

10 The deduced peptide is 759 amino acids long, and has a molecular weight of 88,233. In contrast, N-Glycanase digested, immunopurified FAP $\alpha$  was reported to have an estimated  $M_r$  of 75,000 on NaDodSO<sub>4</sub>/PAGE (Rettig et al., Canc. Res. 53: 3327-3335 (1993)). A TATA box is found 83 base pairs  
15 upstream of the start codon. A polyadenylation signal and a poly-A tail were found in the 5'-untranslated region of the insert.

A GenBank data base search was then carried out. The most closely related genes found were those encoding  
20 dipeptidyl peptidase IV homologues (DPPIV; EC 3.4.14.5), with human DPPIV (also known as T-cell activation antigen CD26), showing 61% nucleotide sequence identity, and 48% amino acid sequence identity.

The second set of related genes are human, rat, and  
25 bovine homologues of DPPX, a gene of unknown function widely expressed in brain and other normal tissues. The predicted human DPPX gene product shows about 30% amino acid sequence identity with FAP $\alpha$  and CD26. The FAP $\alpha$  molecule exhibits structural features typical of type II integral membrane  
30 proteins, including a large COOH-terminal extracellular domain, a hydrophobic transmembrane segment, and a short cytoplasmic tail. The putative extracellular domain contains six potential N-glycosylation sites, 13 cysteine residues (8 of which are conserved between FAP $\alpha$  and CD26), and three  
35 segments corresponding to highly conserved catalytic domains characteristic of serine proteases, such as DPPIV. These conserved sequences are presented in Table 2, which follows. Comparisons to DPPIV and DPPX were made via Morimoto et al.,  
40 supra; Wada et al., Proc. Natl. Acad. Sci. USA 89: 197-201 (1992); Yokotani et al., Human Mol. Genet. 2: 1037-1039 (1993).

5 Table 2. Putative catalytic domains of FAP $\alpha$ , DPPIV and DPPX.

	625	701	733
Human FAP $\alpha$	....WGWSYEI.....	GTADDNV.....	DQNHGLS....
10 Human DPPIV	....WGWSYGG.....	GTADDNV.....	DEDHCIA....
Mouse DPPIV	....WGWSYGG.....	GTADDNV.....	DEDHGIA....
Rat DPPIV	....WGWSYGG.....	GTADDNV.....	DEDHGIA....
Yeast DPPIV	....FGWSYGG.....	GTGDDNV.....	DSDHSIR....
15 Human DPPX	....FGKDYGG.....	PTADEKI.....	DESHYFT....
Rat DPPX	....FGKDYGG.....	ATADEKI.....	DESHYFH....
Bovine DPPX	....FGKDYGG.....	ATEDEKI.....	DESHYFS....

Example 6

20 An additional set of experiments were carried out to determine whether FAP $\alpha$  related sequences are present in non-human species. To do so, human, mouse, and Chinese hamster genomic DNA was digested using restriction enzymes, and tested, via Southern blotting, using the 2.3 kb fragment, labelled with <sup>32</sup>P, describes supra. Hybridization was carried out using stringent washing conditions (0.1 x SSC, 0.1% NaDodSO<sub>4</sub>, 68°C). Cross-hybridization was readily observed with both the mouse and hamster DNA, suggesting the existence of highly conserved FAP $\alpha$  homologues. In control experiments using the CD26 cDNA fragment described supra, no evidence of cross hybridization was observed.

Example 7

35 The CD26 molecule shares a number of biochemical and serological properties with FAP $\beta$ , which is a previously described, FAP $\alpha$  associated molecule having a molecular weight of 105 kd, and found on cultured fibroblasts and melanocytes (Rettig et al., Canc. Res. 53: 3327-3335 (1993)). Cotransfection experiments were carried out to determine whether FAP $\beta$  is a CD26 gene product. To test this, the same protocols were used which were used for transfection with pFAP.38 or pCD26, as described supra, but using the two

5 vectors. The results presented supra showed that cotransfection efficiency was about 40% for each vector, so about 10-20% of cell should be cotransfected.

Following cotransfection, the COS-1 cells were Trans <sup>35</sup>S-labeled, as described supra, then lysed, also as described  
10 supra.

The resulting cell extracts were separated on Con A SEPHAROSE, and the antigen (FAP $\alpha$  and/or CD26) were recovered in the Con A-bound fraction. The bound fraction was eluted with 0.25 M  $\alpha$ -D-mannopyranoside. Immunoprecipitation was then  
15 carried out, as described supra, and the precipitates were separated on NaDodSO<sub>4</sub>/PAGE, also as discussed supra.

Figure 4 shows these results, together with results from single transfection experiments. Those cells transfected only with pFAP.38 produce FAP $\alpha$ , but not FAP $\beta$  (determined from mAb F19 immunoprecipitates). They also produce no CD26 antigen (tested with EF-1). Those cells transfected with pCD26 alone produce CD26 but no FAP $\alpha$ . Cotransfectants produce CD26 and FAP $\alpha$ /FAP $\beta$  heteromers, as determined in the mAb F19 precipitates. This result provides direct evidence that FAP $\beta$   
20 is a CD26 gene product.

25

#### Example 8

It has been observed previously that some cultured human cell types coexpress FAP $\alpha$  and CD26, and show FAP $\alpha$ /CD26 heteromer formation. In vivo distribution patterns of FAP $\alpha$  and CD26, however, as determined in previous immunohistochemical studies, appeared to be non-overlapping. (See Rettig et al., Proc. Natl. Acad. Sci. USA 85: 3110-3114 (1988); Garin-Chesa et al., Proc. Natl. Acad. Sci. USA 87: 7235-7329 (1990); Rettig et al., Canc. Res. 53: 3327-3335 (1993); Stein et al., in Knapp et al., eds. Leukocyte typing IV-white cell differentiation antigens, pp 412-415 (Oxford University Press, N.Y. 1989), pp. 412-415; Möbius et al., J. Exp. Immunol. 74: 431-437 (1988)). In view of the potential significance of FAP $\alpha$ /CD26 coassociation, tissue distribution was reexamined, via side by side immunohistochemical staining  
30 of normal tissues and lesional tissues known to contain FAP $\alpha$   
35  
40

5 fibroblasts or FAP $\alpha$  malignant cells.

To test the samples, they were embedded in OCT compound, frozen in isopentane precooled in liquid nitrogen, and stored at -70°C until used. Five micrometer thick sections were cut, mounted on poly-L-lysine coated slides, air dried, and fixed in cold acetone (4°C, for 10 minutes). The sections were then tested with mAbs (10-20 ug/ml), using the well known avidin-biotin immuno-peroxidase method, as described by, e.g., Garin-Chesa et al., J. Histochem. Cytochem. 37: 1767-1776 (1989); Garin-Chesa et al., Proc. Natl. Acad. Sci. USA 87: 7235-7239 (1990); Rettig et al., Canc. Res. 53: 3327-3335 (1993); Garin-Chesa et al., Am. J. Pathol. 142: 557-567.

The results are shown in figure 5. Breast, colorectal, pancreas and lung carcinomas showed strong expression of FAP $\alpha$  and not CD26 was found (see figures 5A and 5B). Five FAP $\alpha$  sarcomas, including malignant fibrous histocytoma (figures 5C and 5D), were tested, and there was no expression of CD26. Examination of reactive fibroblasts of healing dermal wounds (figures 5E, 5F), showed abundant expression of both FAP $\alpha$  and CD26. The three renal carcinomas tested (figures 5G, 5H), showed expression of CD26 in malignant epithelium. FAP $\alpha$  was absent from malignant epithelial cells, and showed low expression in the stroma of these carcinomas.

The foregoing examples describe an isolated nucleic acid molecule which codes for fibroblast activating protein alpha ("FAP $\alpha$ "). The expression product of the sequence is a protein which, on SDS-PAGE, shows a molecular weight of about 75 kd. Deduced amino acid sequence, as provided in SEQ ID NO: 1, for one form of the molecule, yields a molecular weight of about 88 kd. It is to be understood that, as described, FAP $\alpha$  may be glycosylated, with the type and amount of glycosylation varying, depending upon the type of cell expressing the molecule. The experiment described herein shows this.

The invention also comprehends the production of expression vectors useful in producing the FAP $\alpha$  molecule. In their broadest aspect, these vectors comprise a FAP $\alpha$  coding sequence, operably linked to a promoter. Additional elements

5 may be a part of the expression vector, such as genes which confer antibiotic resistance, amplifiable genes, and so forth.

The coding sequences and vectors may also be used to prepare cell lines, wherein the coding sequence or expression vector is used to transfect or to transform a recipient host.  
10 The type of cell used may be prokaryotic, such as *E. coli*, or eukaryotes, such as yeast, CHO, COS, or other cell types.

The identification of nucleic acid molecules such as that set forth in SEQ ID NO: 1 also enables the artisan to identify and to isolate those nucleic acid molecules which hybridize to it under stringent conditions. "Stringent condition" as used  
15 herein, refers to those parameters set forth supra, whereby both murine and hamster sequences were also identified. It will be recognized by the skilled artisan that these conditions afford a degree of stringency which can be achieved using parameters which vary from those recited. Such variance  
20 is apprehended by the expression "stringent conditions".

The ability of nucleic acid molecules to hybridize to complementary molecules also enables the artisan to identify cells which express FAP $\alpha$ , via the use of a nucleic acid  
25 hybridization assay. One may use the sequences described in the invention to hybridize to complementary sequences, and thus identify them. In this way, one can target mRNA, e.g., which is present in any cell expressing the FAP $\alpha$  molecule.

It is of course understood that the nucleic acid  
30 molecules of the invention are also useful in the production of recombinant FAP $\alpha$ . The recombinant protein may be used, e.g., as a source of an immunogen for generation of antibodies akin to known mAb F19, and with the same uses. Similarly, the recombinant protein, and/or cells which express the molecule  
35 on its surface, may be used in assays to determine antagonists, agonists, or other molecules which interact with the FAP $\alpha$  molecule. Such molecules may be, but are not necessarily limited to, substrates, inhibiting molecules, antibodies, and so forth. This last feature of the invention  
40 should be considered in light of the observed structural resemblances to membrane bound enzymes. This type of molecule

5 is associated with certain properties which need not be described in detail here. It will suffice to say that inhibition or potentiation of these properties as associated with FAP $\alpha$  is a feature of this invention. For example, one may identify substrates or the substrate for the FAP $\alpha$  molecule, via the use of recombinant cells or recombinant FAP $\alpha$  per se. The substrates can be modified to improve their effect, to lessen their effect, or simply to label them with detectable signals so that they can be used, e.g., to identify cells which express FAP $\alpha$ . Study of the interaction of substrate and FAP $\alpha$ , as well as that between FAP $\alpha$  and any molecule whatsoever, can be used to develop and/or to identify agonists and antagonists of the FAP $\alpha$  molecule.

Other aspects of the invention will be clear to the skilled artisan, and need not be set forth here.

20 The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, it being recognized that various modifications are possible within the scope of the invention.



## 5 (1) GENERAL INFORMATION:

(i) APPLICANTS: Rettig, Wolfgang J.; Scanlan, Matthew J.;  
Garin-Chesa, Pilar; Old, Lloyd J.

10 (ii) TITLE OF INVENTION: ISOLATED NUCLEIC ACID MOLECULE CODING  
FOR FIBROBLAST ACTIVATION PROTEIN  $\alpha$  AND USES THEREOF

(iii) NUMBER OF SEQUENCES: 1

15 (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Felte & Lynch

(B) STREET: 805 Third Avenue

(C) CITY: New York City

(D) STATE: New York

20 (E) COUNTRY: USA

(F) ZIP: 10022

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette, 5.25 inch, 360 kb storage

25 (B) COMPUTER: IBM PS/2

(C) OPERATING SYSTEM: PC-DOS

(D) SOFTWARE: Wordperfect

(vi) CURRENT APPLICATION DATA:

30 (A) APPLICATION NUMBER: 08/230,491

(B) FILING DATE: 20-APRIL-1994

(C) CLASSIFICATION: 435

(viii) ATTORNEY/AGENT INFORMATION:

35 (A) NAME: Hanson, Norman D.

(B) REGISTRATION NUMBER: 30,946

(C) REFERENCE/DOCKET NUMBER: LUD 330

(ix) TELECOMMUNICATION INFORMATION:

40 (A) TELEPHONE: (212) 688-9200

(B) TELEFAX: (212) 838-3884

15

5 (2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2812 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

10 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:


AAGAACGCCC CCAAATCTG TTTCTAATTT TACAGAAATC TTTTGAAACT TGGCACGGTA 60  
TTCAAAAGTC CGTGGAAAGA AAAAAACCTT GTCCTGGCTT CAGCTTCCAA CTACAAAGAC 120  
15 AGACTTGGTC CTTTCAACG GTTTTCACAG ATCCAGTGAC CCACGCTCTG AAGACAGAAT 180  
TAGCTAACTT TCAAAAACAT CTGGAAAAAT GAAGACTTGG GTAAAAATCG TATTTGGAGT 240  
TGCCACCTCT GCTGTGCTTG CCTTATTGGT GATGTGCATT GTCTTACGCC CTTCAAGAGT 300  
TCATAACTCT GAAGAAAATA CAATGAGAGC ACTCACACTG AAGGATATTT TAAATGGAAC 360  
ATTTTCTTAT AAAACATTTT TTCCAAACTG GATTTTCAGGA CAAGAATATC TTCATCAATC 420  
20 TGCAGATAAC AATATAGTAC TTTATAATAT TGAAACAGGA CAATCATATA CCATTTTGAG 480  
TAATAGAACC ATGAAAAGTG TGAATGCTTC AAATTACGGC TTATCACCTG ATCGGCAATT 540  
TGTATATCTA GAAAGTGATT ATTCAAAGCT TTGGAGATAC TCTTACACAG CAACATATTA 600  
CATCTATGAC CTTAGCAATG GAGAATTTGT AAGAGGAAAT GAGCTTCCTC GTCCAATTCA 660  
GTATTTATGC TGGTCGCTTG TTGGGAGTAA ATTAGCATAT GTCTATCAA ACAATATCTA 720  
25 TTTGAAACAA AGACCAGGAG ATCCACCTTT TCAATAACA TTTAATGGAA GAGAAAATAA 780  
AATATTTAAT GGAATCCCAG ACTGGGTTTA TGAAGAGGAA ATGCTTCCTA CAAAATATGC 840  
TCTCTGGTGG TCTCCTAATG GAAAATTTT GGCATATGCG GAATTTAATG ATAAGGATAT 900  
ACCAGTTATT GCCTATTCCT ATTATGGCGA TGAACAATAT CCTAGAACAA TAAATATTCC 960  
ATACCCAAAG GCTGGAGCTA AGAATCCCGT TGTTCCGATA TTTATTATCG ATACCACTTA 1020  
30 CCCTGCGTAT GTAGGTCCC AGGAAGTGCC TGTTCAGCA ATGATAGCCT CAAGTGATTA 1080  
TTATTTCACT TGGCTCACGT GGGTTACTGA TGAACGAGTA TGTTTGCACT GGCTAAAAAG 1140  
AGTCCAGAAT GTTTCGGTCC TGTCTATATG TGAATTCAGG GAAGACTGGC AGACATGGGA 1200  
TTGTCCAAAG ACCCAGGAGC ATATAGAAGA AAGCAGAACT GGATGGGCTG GTGGATTCTT 1260  
TGTTTCAAGA CCAGTTTTC GCTATGATGC CATTTCTGAC TACAAAATAT TTAGTGACAA 1320  
35 GGATGGCTAC AAACATATTC ACTATATCAA AGACACTGTG GAAAATGCTA TTCAAATTAC 1380  
AAGTGGCAAG TGGGAGGCCA TAAATATATT CAGAGTAACA CAGGATTCAC TGTTTTATTC 1440  
TAGCAATGAA TTTGAAGAAT ACCCTGGAAG AAGAAACATC TACAGAATTA GCATTGGAAG 1500  
CTATCCTCCA AGCAAGAAGT GTGTTACTTG CCATCTAAGG AAAGAAAGGT GCCAATATTA 1560  
CACAGCAAGT TTCAGCGACT ACGCCAAGTA CTATGCACTT GTCTGCTACG GCCCAGGCAT 1620  
40 CCCCATTTCC ACCCTTCATG ATGGACGCAC TGATCAAGAA ATTAAATCC TGAAGAAAA 1680  
CAAGGAATTG GAAAATGCTT TGAAAAATAT CCAGCTGCCT AAAGAGGAAA TTAAGAAACT 1740

5 TGAAGTAGAT GAAATTACTT TATGGTACAA GATGATTCTT CCTCCTCAAT TTGACAGATC 1800  
AAAGAAGTAT CCCTTGCTAA TTCAAGTGTA TGGTGGTCCC TGCAGTCAGA GTGTAAGGTC 1860  
TGTATTTGCT GTTAATTGGA TATCTTATCT TGCAAGTAAG GAAGGGATGG TCATTGCCTT 1920  
GGTGGATGGT CGAGGAACAG CTTTCCAAGG TGACAACTC CTCTATGCAG TGTATCGAAA 1980  
GCTGGGTGTT TATGAAGTTG AAGACCAGAT TACAGCTGTC AGAAAATTCA TAGAAATGGG 2040  
10 TTTTATTGAT GAAAAAAGAA TAGCCATATG GGGCTGGTCC TATGAGATAC GTTTCATCAC 2100  
TGGCCCTTGC ATCTGGAAC TGTCTTTTCA AATGTGGTAT AGCAGTGGCT CCAGTCTCCA 2160  
GCTGGGAATA TTACGCGTCT GTCTACACAG AGAGATTCAT GGGTCTCCCA ACAAAGATGA 2220  
TAATCTTGAG CACTATAAGA ATTCAACTGT GATGGCAAGA GCAGAATATT TCAGAAATGT 2280  
AGACTATCTT CTCATCCACG GAACAGCAGA TGATAATGTG CACTTTCAA ACTCAGCACA 2340  
15 GATTGCTAAA GCTCTGGTTA ATGCACAAGT GGATTTCCAG GCAATGTGGT ACTCTGACCA 2400  
GAACCACGGC TTATCCGGCC TGTCCACGAA CCACTTATAC ACCCACATGA CCCACTTCCT 2460  
AAAGCAGTGT TTCTCTTTGT CAGACTAAAA ACGATGCAGA TGCAAGCCTG TATCAGAATC 2520  
TGAAAACCTT ATATAAACCC CTCAGACAGT TTGCTTATTT TATTTTTTAT GTTGTAATAAT 2580  
GCTAGTATAA ACAAACAAAT TAATGTTGTT CTAAAGGCTG TTAACAAAAA GATGAGGACT 2640  
20 CAGAAGTTCA AGCTAAATAT TGTTTACATT TTCTGGTACT CTGTGAAAGA AGAGAAAAGG 2700  
GAGTCATGCA TTTTGCTTTG GACACAGTGT TTTATCACCT GTTCATTGTA AGAAAAATAA 2760  
TAAAGTCAGA AGTTCAAAAA AAAAAAAAAA AAAAAAAAAA GCGGCCGCTC GA 2812


5 We claim:

1. Isolated nucleic acid molecule which codes for mammalian FAP $\alpha$  having a molecular weight of about 88 kilodaltons based upon its deduced amino acid sequence.
- 10 2. The isolated nucleic acid molecule of claim 1, wherein said FAP $\alpha$  consists of the amino acid sequence set forth in SEQ ID NO: 1.
- 15 3. The isolated nucleic acid molecule of claim 1, consisting of the nucleotide sequence of SEQ ID NO: 1.
- 20 4. Isolated nucleic acid molecule which hybridizes to the nucleotide sequence of SEQ ID NO: 1, under stringent conditions.
5. Expression vector comprising the isolated nucleic acid molecule of claim 1, operably linked to a promoter.
- 25 6. Cell line transformed or transfected by the isolated nucleic acid molecule of claim 1.
7. Cell line transformed or transfected by the expression vector of claim 5.
- 30 8. Method for determining expression of FAP $\alpha$  in a cell comprising contacting said cell with the isolated nucleic acid molecule of claim 1 and determining hybridization of said isolated nucleic acid molecule to a complementary sequence in said cell as a determination of expression of FAP $\alpha$ .
- 35


FIG. 1



**FIG. 2A**



**FIG. 2B**



3 / 5

## FIG. 3

FAP	1	MKTWVKIVFGV*ATSAVLALLVMCIVLRPSRVHNSEENTMRALTLKDILN	49
CD26	1	---PW-VLL-LLGAA-LVTIITVPV--LNKGTDDATADSRKTY--T-Y-K	50
FAP	50	GTFSYKTFPPNWISGQEYLHQSadNNIVLYNIETGQSYTILSNRTMKSV*	98
CD26	51	N-YRL-LYSLR---DH---YKQ*E---LVF-A-Y-N-SVF-E-S-FDEFG	99
FAP	99	*NASNYGLSPDRQFVYLESDYSLWRYSYTATYIYDLSNGEFVRGNELP	147
CD26	100	HSIND-SI---G--IL--YN-V-Q--H----S-D----NKRQLITEERI-	149
		<u>fap-1</u>	
FAP	148	RPIQYLCWSPVGSKLAYVYQNNIYLKQRPDPPFQITFNGRENKIFNGIP	197
CD26	150	NNT-WVT-----H-----WN-D--V-IE-NL-SYR--WT-K-DI-Y---T	199
		<u>fap-2</u>	
FAP	198	DWVYEEEMLPKYALWWSPNGKFLAYAEFNDKDIPVIAYSYYGDE**QYP	245
CD26	200	-----VFSAYS-----T-----Q---TEV-L-E--F-S--SL---	249
FAP	246	RTINIPYPKAGAKNPVVRIFIIDT***TYPAYVGPQEVVPAMIASSDY	292
CD26	250	K-VRV-----V--T-KF-VVN-DSLSSVTNATSIQITA--SMLIG-H-	299
FAP	293	FSWLTWVTDERVCLQWLKRVONVSVLSICDFREDWQTWDCPKTQEHIEES	342
CD26	300	LCDV--A-Q--IS----R-I--Y--MD---YD-SSGR-N-LVARQ---M-	349
FAP	343	RTGWAGGFFVSRPVFSYDAISYYKIFSDKDGKXHIHYIKDTVENAIQITS	392
CD26	350	T---V-R-RP-E-H-TL-GN-F---I-NEE--R--C-FQIDKKDCTF--K	399
FAP	393	GKWEAINIFRVTQDSLIFYSSNEFEYYPGRRNIRISIGSYPPSKKCVTCH	442
CD26	400	-T--V-G-EAL-S-Y-Y-I---YKGM--G--L-K-QLSD-T*KVT-LS-E	448
FAP	443	LRKERCQYYTASFSDYAKYYALVCYGPPISTLHDGRTDQEIKILEENK	492
CD26	449	-NP-----SV---KE----Q-R-S---L-LY---SSVN-KGLRV--D-S	498
		<u>fap-3</u>	
FAP	493	ELENALKNIQLPKEEIKLEVDEITLWYKMILPPQFDRSKKYPLLIQVYG	542
CD26	499	A-DKM-Q-V-M-SKKLDFIILN-TKF--Q-----H--K-----LD--A	548
FAP	543	GPCSQSVRSVFAVNWISYLASKEGMVIALVDGRGTAFQGDKLLYAVYRKL	592
CD26	549	-----KADT--RL--AT-----T-NIIV-SF-----SGY-----IMH-IN-R-	598
FAP	593	GVYEVEDQITAVRKFIEMGFIDEKRIAIWGSYEIRFITGPCIWNWSFQM	642
CD26	599	-TF-----E-A-Q-SK---V-N-----GGYVTSMVLGSGSVGFK	648
FAP	643	WYSSGSSLQLGILRVCLHRE*IHGSPNKDDNLEHYKNSTVMARAEYFRNV	691
CD26	649	CGIAVAPVSRWEYYSVYT-RYM-L-TPE---D--R-----S---N-KQ-	698
FAP	692	DYLLIHGTADDNVHFQNSAQIAKALVNAQVDFQAMWYSDQNHGLSGLSTN	741
CD26	699	E-----Q-----S-----DVG-----T-ED--IASSTAH	748
FAP	742	*HLYTHMTHFLKQCFSLS	
CD26	749	Q-I----S--I-----P	

4/5


Fig. 5

Breast cancer	MFH	Healing wound	Renal cancer	
⊕ A	⊕ C	⊕ E	⊖ G	FAP $\alpha$
⊖ B	⊖ D	⊕ F	⊕ H	CD26

Immunohistochemistry (see Kodachromes)



Fig. 4



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/04860

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 15/00, 15/09, 15/12

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 69.1, 172.3, 240.2, 252.3, 320.1; 536/23.5, 24.3, 24.31, 24.33

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS and DIALOG (files 5, 155,351,357,358) search terms: FAP, fibroblast activation protein, antigen, CD26

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y ---- A	CANCER RESEARCH, Volume 53, issued 15 July 1993, W.J. Rettig et al, "Regulation and Heteromeric Structure of the Fibroblast Activation Protein in Normal and Transformed Cells of Mesenchymal and Neuroectodermal Origin", pages 3327-3335, see entire document.	1-7 ----- 8
Y ---- A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES USA, Volume 85, issued May 1988, W.J. Rettig et al, "Cell-surface glycoproteins of human sarcomas: Differential expression in normal and malignant tissues and cultured cells", pages 3110-3114, see entire document.	1-7 ----- 8

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A		document defining the general state of the art which is not considered to be of particular relevance
*E		earlier document published on or after the international filing date
*L		document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
*O		document referring to an oral disclosure, use, exhibition or other means
*P		document published prior to the international filing date but later than the priority date claimed
	*X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
	*Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
	*G	document member of the same patent family

Date of the actual completion of the international search

21 JUNE 1995

Date of mailing of the international search report

10 JUL 1995

 Name and mailing address of the ISA/US  
 Commissioner of Patents and Trademarks  
 Box PCT  
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

MARIANNE PORTA ALLEN

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/04860

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES USA, Volume 84, issued December 1987, A. Aruffo et al, "Molecular cloning of a CD28 cDNA by a high-efficiency COS cell expression system", pages 8573-8577, see entire document.	1-7

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/04860

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/6, 69.1, 172.3, 240.2, 252.3, 320.1; 536/23.5, 24.3, 24.31, 24.33